



# SENCOR® 75

## Turf Herbicide

## Specimen Label

*For control of certain broadleaf and grass weeds in turf.*

**ACTIVE INGREDIENT:**

Metribuzin

4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one . . . . . 75%

**INERT INGREDIENTS:** . . . . . 25%

Total: 100%

Store in a cool, dry place.

EPA Reg. No. 264-738

**STOP - Read the Label Before Use.  
KEEP OUT OF REACH OF CHILDREN.**

### CAUTION

**PRECAUCION AL USUARIO:** Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

**PERSONAL PROTECTIVE EQUIPMENT:**

**Applicators and Other Handlers Must Wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made out of any water-proof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

NET CONTENTS:

**BACKED  
by BAYER™**

### User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Do not contaminate feed or food. Keep out of reach of children.

Obtain prompt medical aid if poisoning should occur.

**Symptoms of Poisoning:** The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by breathing difficulties and sedation.

### FIRST AID

|                               |   |
|-------------------------------|---|
| <b>If swallowed</b>           | <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul> |
| <b>If on skin or clothing</b> | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>  |
| <b>If in eyes</b>             | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>  |

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a SENCOR container or label with you when calling a poison control center or doctor, or going for treatment.

**Note To Physician:** Treat the patient symptomatically.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**Ground Water Advisory:** Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply metribuzin where the water table (ground water) is close to the surface and where the soils are very permeable, i.e., well drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## GENERAL INFORMATION

### GENERAL PRECAUTIONS AND RESTRICTIONS

Do not enter or allow others to enter treated area until sprays have dried. Applications should not be made to dormant turf in the transitional bermudagrass growing zones which are or can be expected to be adversely affected by cold weather stress.

Low-pressure and high-volume hand-wand equipment is prohibited.

Apply no more than 2 lbs SENCOR 75 Turf Herbicide (1.5 lbs active ingredient) per acre in a single growing season. Apply no more than 1 time to dormant turf and 2 times to actively growing turf in a single growing season.

Avoid spray overlaps that will increase dosages above those recommended.

Do not apply aerially.

Apply only to established bermudagrass turf which has a mowing height of one-half inch or greater. Do not apply to greens, tees, aprons, or other areas of closely mowed turf.

Do not apply within the root zone area of ornamentals, shrubs, or trees as phytotoxicity may occur.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Delay mowing treated areas for at least 3 days for maximum weed control. For best results, delay mowing until after rainfall or irrigation is received. Grass clippings cannot be used for animal feed.

When applying SENCOR to actively growing turf, use the lower rate in areas where soil pH is greater than 7.5.

Do not use on crops grown for food or forage.

Do not allow sprays to drift on to adjacent desirable plants.

**NOTE: Not for use in commercial greenhouses, nurseries, on sod farms or on grass grown for seed. For use on plants intended for aesthetic purposes or climatic modification and being grown on golf courses or lawns and grounds.**

Observe all cautions and limitations on labeling of all products used in mixtures.

### MIXING

Fill the spray tank 1/4 to 1/3 full with clean water and add SENCOR 75 Turf Herbicide at the recommended rate. Mix thoroughly and add water to fill spray tank. Maintain sufficient agitation during mixing and spraying operations to insure a uniform spray mixture.

It is essential that the sprayer be accurately calibrated before applying SENCOR 75 Turf Herbicide. Avoid boom-overlaps that will increase dosages above those recommended. Check the sprayer frequently during application to be sure it is working properly and delivering a uniform spray pattern.

Do not apply this product through any type of irrigation system.

### SPRAYER CLEANUP

Spray equipment must be thoroughly cleaned to remove remaining traces of herbicide that might injure other areas to be sprayed. Drain any remaining spray solution of SENCOR from the spray tank and discard in non-crop areas away from water supplies. Rinse the spray tank and refill with water, adding a heavy-duty detergent at the rate of one cup per 20 gallons of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of spray tank, nozzles or spray rig.

**BERMUDAGRASS TURF (Established):**

SENCOR 75 Turf Herbicide is recommended for application by commercial applicators only on established bermudagrass turf (such as parks, athletic fields, golf course fairways and cemeteries) which has a mowing height of one-half inch or greater. (See “General Precautions and Restrictions” for area of use).

| <b>WEEDS CONTROLLED (Dormant Turf)</b>  |   |
|---|---|
| SENCOR 75 Turf Herbicide, applied according to directions when weeds are present and actively growing in bermudagrass turf, will effectively control:                                       |   |
| <b>BROADLEAF WEEDS (EXCEPT CALIFORNIA)</b>  |   |
| Alkali mallow (a. sida)<br>Bedstraw<br>Carolina geranium<br>Carpetweed<br>Common chickweed<br>Corn speedwell<br>Henbit<br>Hop clover<br>Spotted bur clover<br>White clover<br>London rocket | Nettleleaf goosefoot<br>Parsley-piert<br>Prostrate knotweed<br>Red deadnettle<br>Silversheath knotweed<br>Shepherdspurse<br>Small flowered buttercup<br>Spotted spurge<br>Spur weed<br>Wild mustard |
| <b>BROADLEAF WEEDS (CALIFORNIA ONLY)</b>  |   |
| Alkali mallow (a. sida)<br>Carpetweed<br>Common chickweed<br>London rocket  | Nettleleaf goosefoot<br>Shepherdspurse<br>Wild mustard  |

| <b>WEEDS CONTROLLED (Actively Growing Turf)</b>  |   |
|--|---|
| SENCOR 75 Turf Herbicide, applied to actively growing bermudagrass turf according to directions, will effectively control: |   |
| <b>GRASS WEEDS</b>   |   |
| Annual bluegrass<br>(Poa annua)<br>Goosegrass<br>(Except California)   | Littleseed canarygrass<br>Rabbitfootgrass |

| <b>RECOMMENDED APPLICATIONS</b>   |  |
|---|--|
| <b>ESTABLISHED BERMUDAGRASS TURF</b><br>(which has a mowing height of one-half inch or greater) |  |
| POUNDS<br>SENCOR 75 TURF<br>HERBICIDE   | Remarks  |
| 2/3   | <b>DORMANT TURF:</b> Apply specified dosage in 40 gallons of water per acre as a uniform broadcast spray prior to green up of turf. See “General Precautions and Restrictions” prior to using this product on turf.  |
| 1/3 to 2/3  | <b>ACTIVELY GROWING TURF:</b> Apply specified dosage in 40 gallons of water per acre as a uniform broadcast spray. Application should be made only when turf is vigorously growing and not under stress conditions. Repeat if necessary, but do not apply closer than 1 week intervals. Do not make more than 2 applications to actively growing turf per season. Applications may result in temporary discoloration which turf soon outgrows. See “General Precautions and Restrictions” prior to using this product on turf. |

**Other Weeds Controlled:** SENCOR 75 Turf Herbicide, tank mixed with MSMA and applied to actively growing bermudagrass turf according to directions will effectively control:

- Barnyardgrass
- Common yellow woodsorrel
- Crabgrass
- Dallisgrass
- Nutsedge
- Sandbur

For control of these weeds, apply SENCOR as directed above and use as a tank mix with MSMA. Consult the MSMA label or contact your local turf extension specialist for additional directions, rates, weed species controlled, and precautions.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
©2003 Bayer CropScience

010806A

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Bayer CropScience's control it is impossible for Bayer CropScience to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES, RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Bayer AG, 2003