

# Drive<sup>®</sup> 75 DF

herbicide

**Active ingredient**

3,7-dichloro-8-quinolinecarboxylic acid.....	75.0%
Inert Ingredients.....	25.0%
<b>Total</b> .....	<b>100.0%</b>

**KEEP OUT OF REACH OF CHILDREN.**  
**CAUTION**

EPA Registration Number 7969-130

**Precautionary Statements**

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Causes moderate eye injury. May cause allergic skin response.

---

**Directions For Use**


---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

All applicable **Directions, Restrictions and Precautions** are to be followed. This labeling must be in the user's possession during application.

---

**General Restrictions and Limitations**


---

- Do not apply more than 2 pounds of **Drive<sup>®</sup> 75 DF herbicide** per acre (or 0.73 ounces per 1,000 square feet) in one year.
- Do not apply to fine fescue unless it is part of a seed blend.
- Do not apply to bahiagrass, carpetgrass, St. Augustinegrass, centipedegrass, dichondra, or lawns or turf where desirable clovers are present.
- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass and perennial ryegrass.
- Do not apply to exposed feeder roots of trees or ornamentals. Be particularly careful within the drip line of trees and other ornamental species.
- Do not apply into any ornamental bed.
- Do not use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Apply when wind speed is less than 10 mph as drift may cause damage or death of non-target area vegetation. Do not apply when conditions favor drift from target area.
- Use a lawn type sprayer with coarse spray as wind drift is less likely.
- Avoid mist and spray onto vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants, **especially plants belonging to the Solanaceae family such as tomatoes, eggplants, and bell peppers.**
- Do not pour spray solutions near these plants.
- Do not apply to golf course collars or greens.
- Do not apply through any type of irrigation system.
- Do not use to formulate or reformulate any other pesticide product which is not registered by EPA.
- For best results, weeds should not be under stress from lack of water, excessive water, low fertility, mowing shock, excessive hot or cold temperatures, or injury from other herbicide applications.

# BASF

## Application Rates and Timing for Postemergence Weed Control in Turf.

Weed	Drive 75 DF Rate	Additive Rate
Barnyardgrass Bindweed, Field Black medic Clover, Hop , Red , White Crabgrass, Large <sup>2,5</sup> , Smooth <sup>2,5</sup> Dandelion, Common <sup>3</sup> Dollarweed Foxtails, Giant <sup>2</sup> , Green <sup>2</sup> , Yellow <sup>2</sup> Kikuyugrass <sup>1</sup> Signalgrass, Broadleaf <sup>2</sup> Speedwell, Common , Slender , Thymeleaf Torpedograss <sup>4</sup>	1 pound per acre or 0.367 ounce per 1,000 square feet	1.5 pints per acre (0.55 oz per 1,000 square feet) methylated seed oil
<sup>1</sup> Make 2 sequential applications of 0.367 ounce of <b>Drive 75 DF</b> per 1,000 square feet or 3 sequential applications of 0.245 ounce of Drive 75 DF per 1,000 square feet at 14 day intervals. <sup>2</sup> Apply either before second tiller or as weed grasses mature. <sup>3</sup> Tank mix partner or sequential application required. <sup>4</sup> Make 2 sequential applications of 0.367 ounce of <b>Drive 75 DF</b> per 1,000 square feet or 3 sequential applications of 0.245 ounce of Drive 75 DF per 1,000 square feet at 21 day intervals. <sup>5</sup> Not for use in California.		

### Seeding / Overseeding / Sprigging Timing Chart

Variety	7 days before seeding	At Seeding <sup>1</sup>	7 days after emergence	14 days after emergence	28 days after emergence
Annual Bluegrass	OK	OK	OK	OK	OK
Annual Ryegrass	OK	OK	OK	OK	OK
Buffalograss	OK	OK	OK	OK	OK
Common Bermuda	OK	OK	OK	OK	OK
Creeping Bentgrass	OK	NO	NO	NO	OK
Fine Fescue (in blend)	OK	NO	NO	NO	OK
Hybrid Bermuda	OK	OK	OK	OK	OK
Kentucky Bluegrass	OK	NO	NO	NO	OK
Perennial Ryegrass	OK	OK	NO	NO	OK
Tall Fescue	OK	OK	OK	OK	OK
Zoysia	OK	OK	OK	OK	OK

<sup>1</sup> Seeding or Sprigging

Note: No adjuvant or additive should be used when Drive 75 DF applications are made on newly emerged turf seedlings until 28 days after emergence.

Application of Drive 75 DF should be timed around the seeding operations using the above chart as a reference point.

*Drive is a registered trademark of BASF AG.*

©2000 BASF Corporation  
All rights reserved  
NVA 2000-04-073-0050

BASF Corporation  
Agricultural Products  
P.O. Box 13528  
Research Triangle Park, NC 27709